

# A Six-Month Clinical Study to Evaluate the Effects of Sodium Tripolyphosphate and Tetrapotassium Pyrophosphate Based Calculus Dissolution Oral Rinse in Patients with Zirconium Dioxide and Titanium Dental Implants

Vladimir Kokovic<sup>1</sup>, Mustafa Tattan<sup>2</sup>

## ABSTRACT

**Aim:** To clinically evaluate the effects of a novel anti-calculus mouth rinse containing sodium tripolyphosphate and tetrapotassium pyrophosphate (Periogen, USA) on development of gingivitis and plaque around dental implants versus control over a period of six months. **Material and methods:** This was a randomized, 6 month, parallel groups, double blind, single center clinical trial. Forty subjects with present dental implants (22 zirconium dioxide and 20 titanium) were randomly assigned to one of two subgroups: control group (regular brushing) and test group (regular brushing followed by using the calculus dissolution based oral rinse). All subjects were assessed with gingival index (GI), plaque index (PI) and Volpe-Manhold calculus index (VMI) after 3 and 6 months. **Results:** Statistical analysis found that test group in both zirconium dioxide and titanium group demonstrated statistically significant lower GI, PI and VMI scores. **Conclusion:** This study demonstrates that calculus dissolution based Periogen® mouthrinse provided clinically significant reduction in calculus formation in subjects with zirconium dioxide and titanium dental implants when used twice daily for 6 months as an adjunct to toothbrushing.

**Key words:** Dental implant, Oral rinse, Anti-calculus, Peri-implantitis, plaque index, biopolymers, clinical trial

## INTRODUCTION

A dental implant is an artificial tooth root that interfaces with the bone of the jaw or skull through dynamic process of osseointegration to hold a replacement tooth or prosthetic bridge. Dental implants are considered to be safe treatment options for missing tooth or teeth subject to proper case selection and placement technique.

Review of literature on peri-implant soft tissue around zirconia showed that zirconia dioxide implants and abutments provide a very good peri-implant soft tissue interface that achieved an irritation-free attachment. Various *in vivo* and *in vitro* investigations of soft tissue response around zirconia revealed comparable or even better healing response, less inflammatory infiltrate and reduced plaque adhesion on zirconium oxide discs compared to conventionally pure titanium.<sup>1-6</sup>

*In vitro* experimental studies on zirconium showed favorable response in terms of adhesion and proliferation of various cellular components of connective tissue suggesting zirconium as a material of choice for perio-integration.<sup>7</sup> Results of Degidi *et al.* study showed a lower inflammatory response in biopsies

from peri-implant soft tissue around zirconium healing caps when compared to titanium.<sup>8</sup>

The greatest biological challenge that affects implants longevity is poor plaque control and deposition of calculus around dental implants leading to host induced inflammatory response. Dental implants and their components act as a scaffold on which bacteria can accumulate and form bacterial biofilm which can lead to peri-implantitis. It is the most common causes of dental implant failure.<sup>9</sup> It has been concluded that zirconia surface accumulate significantly less bacteria when comparable with titanium ones<sup>10-12</sup>

The progress of science and technology leading to new products development that eradicate the reservoir of bacteria i.e. dental calculus seems to be perfect scientific strategy to add longevity to dental implants and their healthy existence in oral cavity.

Calculus dissolution based oral rinse Periogen®, (Periogen Company USA), presents significantly potential oral rinse for suppressing the mineralization of dental plaque into calculus.<sup>14</sup> A six month comparative study by Saini R proved that Periogen® is clinically proven to prevent tartar/calculus build-up 45% more effectively than brushing alone on new supragingival calculus formation.<sup>13</sup>

**Cite this article :** Kokovic V, Tattan M.A Six-Month Clinical Study to Evaluate the Effects of Sodium Tripolyphosphate and Tetrapotassium Pyrophosphate Based Calculus Dissolution Oral Rinse in Patients with Zirconium Dioxide and Titanium Dental Implants BEMS Reports, 2016;2(2):64-70

## Vladimir Kokovic<sup>1</sup>, Mustafa Tattan<sup>2</sup>

<sup>1</sup>Department of Advance Europe Medical Centre L.L.C, Sharjah UAE  
Maktum Bin Hamdan Dental University College, Dubai UAE

<sup>2</sup>College of Dental Science, Ras al-Khaimah Medical and Health Sciences University, Ras Al Khaimah, UAE

### Correspondence

Vladimir Kokovic

Professor, DDS, MSc, PhD

Advance Europe Medical Centre L.L.C, Sharjah UAE

Maktum Bin Hamdan Dental University College, Dubai UAE

Email: aemcuae@gmail.com

Ph.no: +971 65574480

### History

- Submission Date: 26-10-16;
- Review completed: 18-11-16;
- Accepted Date: 22-11-16.

DOI : 10.5530/BEMS.2016.2.11

### Article Available online

<http://www.bemsreports.org>

### Copyright

© 2016 Phcog.Net. This is an open-access article distributed under the terms of the Creative Commons Attribution 4.0 International license.

In the current literature there is no study of clinical effect of calculus dissolution around dental implants. The aim of this study was to evaluate the efficacy of calculus dissolution based oral rinse (Periogen®) in patients with zirconium dioxide and titanium dental implants placed in lower jaw for long-term effects on prevention of peri-implantitis.

## MATERIALS AND METHODS

The present study was performed in the Dental department of Advance Europe Medical Centre, UAE, after approved from the research and ethical committee of Ras al-Khaimah Medical and Health Sciences University (RAKMHSU), UAE, vide no. 3-2014-FD.

Forty patients present with one or more dental implant in posterior part of lower jaw were analyzed in this study. Twenty patients were treated with zirconium dioxide dental implants (Axis Biodental, Switzerland), polyetheretherketoneketone (PEKK) abutments and full ceramic single crowns on them (Group A), Figure 1-2 and twenty with titanium implants (Starumann, Switzerland), titanium abutments and porcelain-fused to metal single crowns (Group B), Figure 4-5.

The inclusion and exclusion criterion for the subject selection in the study is illustrated in table 1. On the beginning of study patients were asked to understand study protocol and accepted subjects signed the consent form to participate in the study.

## Clinical Evaluation

The subjects were asked to participate for evaluation of following clinical parameters on baseline, 3 and 6 months.

Gingival index (GI)

Plaque index (PI)

Volpe-Manhold calculus index (VMI)

### *Gingival index (GI) (Mombelli)*

In the order to assess the severity of peri-implant soft tissue, the gingival index was used. The peri-implant tissue was divided in four analyzed unites (disto-facial papilla, facial gingival margin, mesio-facial papilla & lingual gingival margin). A score ranging from 0 to 3 was given to each surface using the periodontal probe. Score of the all unites were summed and divided by four yielded the individual implant score. Totalling all the scores per implant and dividing by implants number was used for calculation of GI per person.

### *Plaque index (PI) (Mombelli)*

Plaque was assessed on the four peri-implant surfaces (mesial, distal, buccal and lingual) with score between 0 to 3. Plaque score for each implant was obtained by totaling the score for each implant and individual score was calculated.

**Table 1: Inclusion and Exclusion Criteria for the Subjects**

Inclusion criteria:	
•	Age-group between 18 and 65
•	ASA I or II group patient
•	Patients with present one or more dental implants (zirconium dioxide or titanium) in lower jaw with single crown restoration.
Exclusion criteria:	
•	Presence of any neurological disorders
•	Patient who received antibiotics or NSAIDS in the past 9 to 11 weeks
•	Pregnant and lactating mothers
•	Patients with clinically mobile implants
•	Heavy smokers
•	The oral hygiene is inadequate

**Table 2: Distribution of mean and standard deviation (SD) values of clinical parameters in Group A (Test and Control)**

Clinical Parameters	Base Level Mean $\pm$ SD	3 months Mean $\pm$ SD	6 months Mean $\pm$ SD	Percentage Decrease
Test Group				
GI	0.49 $\pm$ 0.13	0.37 $\pm$ 0.10	0.18 $\pm$ 0.08	63 %
PI	0.98 $\pm$ 0.39	0.72 $\pm$ 0.18	0.49 $\pm$ 0.11	50%
VMI	1.93 $\pm$ 0.69	1.42 $\pm$ 0.72	0.93 $\pm$ 0.91	51%
Control Group				
GI	0.57 $\pm$ 0.27	0.50 $\pm$ 0.19	0.42 $\pm$ 0.16	26%
PI	1.01 $\pm$ 0.42	0.97 $\pm$ 0.39	0.89 $\pm$ 0.41	11%
VMI	2.03 $\pm$ 0.91	1.87 $\pm$ 0.63	1.78 $\pm$ 0.45	12%

**Table 3: Distribution of mean and standard deviation (SD) values of clinical parameters in Group B (Test and Control)**

Clinical Parameters	Base Level Mean ± SD	3 months Mean ± SD	6 months Mean ± SD	Percentage Decrease
Test Group				
GI	0.91 ± 0.31	0.73 ± 0.47	0.44 ± 0.27	51%
PI	1.3 ± 0.51	0.87 ± 0.67	0.63 ± 0.31	51%
VMI	2.91 ± 1.1	1.79 ± 0.83	1.53 ± 0.97	47%
Control Group				
GI	1.01 ± 0.47	0.97 ± 0.4	0.87 ± 0.51	14%
PI	1.19 ± 0.43	1.12 ± 0.51	1.02 ± 0.42	6%
VMI	3.18 ± 1.11	2.78 ± 0.92	2.63 ± 0.89	17%



**Figure 1:** Group A - PEKK abutment on zirconium dioxide dental implants



**Figure 2:** Group A - Full ceramic crown



**Figure 3:** Group B - Titanium abutment on titanium implant




**Figure 4:** Group B - Porcelain fused on metal crown




**Figure 5:** Group A - check point at 6 months.




**Figure 6:** Group B - check point at 6 months.




**Graph 1:** Gingival index – Differences between control and test subgroups in zirconium dioxide implants group.




**Graph 2:** Plaque index – Differences between control and test subgroups in zirconium dioxide implants group.




**Graph 3:** Volpe-Manhold calculus index – differences between control and test subgroups in zirconium dioxide implants group.



**Graph 4:** Gingival index – differences between control and test subgroups in titanium implants group.



**Graph 5:** Plaque index – differences between control and test subgroups in titanium implants group.



**Graph 6:** Volpe-Monhold calculus index – differences between control and test subgroups in titanium implants group.

## Volpe-Manhold calculus index (VMI)

To obtain the VMI score, the three tooth planes (mesial, distal and gingival) on lingual surface were exterminated. The periodontal probe was used to measure supragingival calculus on the implant shoulders. The calculus was measured in increment of 0.5mm from 0 to 5mm. Score of the all three planes were summed and all implants scores were summed for subject's total VMI score. The each of forty patients were consider as the statistic unit, that presents statically minimum value for pilot study.

## II Randomization

The randomization was carried out using a lot. The control or test group was randomized using sealed envelopes. Each group (A & B) was divided in to the two subgroups (control & test).

### Control group

Patients from control group were advised to brush twice daily 5 minutes;

### Test group

Patients from test group were advised to brush twice daily 5 minutes followed by using the calculus dissolution based oral rinse Periogen®, USA, (One spoonful of Periogen was dissolved in 100ml of warm water and was directed to rinse with solution for at least one minute in three 20 seconds interval).

## III Statistical Evaluation

All data were first analyzed by descriptive methods (QQ plots, box plots) (SPSS 18.0; SPSS, Austin, TX, USA). The patients were chosen as the unit for statistical analysis. The Wilcoxon signed rank test was used as well as Pearson's coefficient of correlation. The level of significance chosen in all statistical tests was at  $P < 0.05$ .

## RESULTS

40 adult patients (28 females and 12 male) with mean ages 50.37 (SD ± 7.94) were involved in the study. All analyzed implants were tissue-level implants. In group A, 22 two-pieces zirconium dioxide implants (Axis Biodental, Les Bios, Switzerland) with 10 mm length and diameter 4mm were used for single tooth replacement in premolar and molar area of lower jaw. Results of clinical parameters form Group A (zirconium-dioxide dental implant) are presented in Table 1 For all clinical parameters statistical significant differences has been noted between test and control group in the assessment point of 6 months ( $P = 0.02$ ) as illustrated in Figure 5 and Graph 1-3. In group B, 20 titanium SLActive (Straumann, Switzerland) implants length 8mm and 4.1, diameter were used. Results of clinical parameters form Group B (Titanium dental implant) are presented in Table 2. For all clinical parameters statistical significant differences has been noted between test and control group in the check point of 6 months ( $P = 0.009$ ) as illustrated in Figure 6, Graph 4-6.

## DISCUSSION

Similar study evaluated the periodontal parameters with zirconium vs titanium abutments. Less bleeding on probing and less plaque accumulation has been noted around zirconium abutment.<sup>15</sup> Kohal *et al*, compared clinical and radio-graphical outcome of one-piece zirconium implants and natural teeth with fixed dental prosthesis. The modified bleeding index (mBI) and modified plaque index (mPI) were significantly lower at implants than at teeth after one year follow up period.<sup>16</sup>

Results of present study is showing statistically significant better peri-implant condition around soft-tissue zirconium dioxide vs titanium implants in group treated with calculus dissolution based oral rinse Periogen® (Periogen Company-USA). During six months follow-up

period no side effects were recorded among any group using Periogen mouthrinse. Supplementing toothbrushing by Periogen® mouthrinse will lead to success of peri-implant tissue around zirconium and titanium dental implants with significant amount of plaque and calculus reduction.

## CONCLUSION

We conclude that calculus dissolution based Periogen® mouthrinse provided clinically significant reduction in calculus formation in subjects with zirconium dioxide and titanium dental implants when used twice daily for 6 months as an adjunct to toothbrushing.

## ACKNOWLEDGMENT

None

## CONFLICT OF INTEREST

None

## ABBREVIATIONS USED

None

## REFERENCES

- Mellinghoff J. Quality of the peri-implant soft tissue attachment of zirconia implants-abutments. *Z. Zahnärztl. Implants.* 2010;26:62-71.
- Pae A, Lee H, Kim HS, Kwon YD, Woo YH. Attachment and growth behavior of human gingival fibroblasts on titanium and zirconia ceramic surfaces. *Biomed. Mater.* 2009;4(2):025005-12. <https://doi.org/10.1088/1748-6041/4/2/025005> PMID:19208938.
- Rimondini L, Cerroni L, Carrassi A, Torricelli P. Bacterial colonization of zirconia ceramic surfaces: An in vitro and in vivo Study. *Int. J. Oral Maxillofac. Implants.* 2002;17(6):793-8. PMID:12507238.
- Scarano A, Piattelli M, Caputi S, Favero G.A, Piattelli A. Bacterial adhesion on commercially pure titanium and zirconium oxide discs: An in vivo human study. *J. Periodontol.* 2004;75(2):292-6. <https://doi.org/10.1902/jop.2004.75.2.292> PMID:15068118.
- Degidi M, Artese L, Scarano A, Perrotti V, Gehrke P, Piattelli A. Inflammatory infiltrate, microvessel density, nitric oxide synthase expression, vascular endothelial growth factor expression, and proliferative activity in peri-implant soft tissues around titanium and zirconium oxide healing caps. *J. Periodontol.* 2006;77:73-80. <https://doi.org/10.1902/jop.2006.77.1.73> PMID:16579706.
- Größner-Schreiber B, Herzog M, Hedderich J, Dück A, Hannig M, Griepentrog M. Focal adhesion contact formation by fibroblasts cultured on surface-modified dental implants: An in vitro study. *Clin. Oral Implants Res.* 2006;17(6):736-45. <https://doi.org/10.1111/j.1600-0501.2006.01277.x> PMID:17092235.
- Grossen-Schreiber B, Herzog M, Hedderich J, Duck A, Hanning M, Griepentrog M. Focal adhesion contact formation by fibroblasts culture on surface-modified dental implants: an in vitro study. *Clin Oral Implant Res.* 2006;17(6):736-45. <https://doi.org/10.1111/j.1600-0501.2006.01277.x> PMID:17092235.
- Degidi M, Perrotti V, Gehrke P, Piattelli A. Inflammatory infiltrate, microvessel density, nitro oxide synthase expression, vascular endothelial growth factor expression and proliferative activity in perio-implant soft tissue around titanium and zirconium oxide healing caps. *J. Periodontol.* 2006;77(1):73-80. <https://doi.org/10.1902/jop.2006.77.1.73> PMID:16579706.
- Esposito M, Worthington HV. Treatment of peri-implantitis: what interventions are effective? A Cochrane systematic review. *Eur J Oral Implantol.* 2012;5(1):21-41.
- Rimondi LCL, Carrassi A, Torricelli P. Bacterial colonization of zirconia ceramic surface: an in vivo study. *Int. J Oral & Maxilfac. Impl.* 2002;17(6):793-8.
- Scarano MP, Caputi S, Favero GA, Piattelli A. Bacterial Adhesion on Commercially Pure Titanium and Zirconium Oxide Disks: An In vivo Human Study Journal of Periodontology. 2004;75(2):292-6 <https://doi.org/10.1902/jop.2004.75.2.292> PMID:15068118.
- Al-Radhaa DD, Younesb C, O'Sullivan D. Surface properties of titanium and zirconia dental implant materials and their effect on bacterial adhesion. *Journal of Dentistry.* 2011;40(2):146-53. <https://doi.org/10.1016/j.jdent.2011.12.006> PMID:22182466.
- Saini R. Dental calculus: A strategic review. *Int J Dental Health Sci.* 2014;1(5):788-95.
- Saini R. A prospective experimental comparative study on the clinical effects of calculus dissolution based oral rinse in gingival patients. *International Journal of Experimental Dental Science.* 2015;4(1):33-9. <https://doi.org/10.5005/jp-journals-10029-1092>.

15. Van Brakel R, Cune MS; van Winkelhof AJ, de Putter C, Verhoven JW, van de Reijden W. Early bacterial colonization and soft tissue healing around zirconium and titanium abutments: An in vivo study in man. *Clin Oral Implant Res.* 2011;22:571-7. <https://doi.org/10.1111/j.1600-0501.2010.02005.x> PMID:21054554.
16. Kohal RJ, Patzelt SB, Buz F, Shlin H. One-piece zirconium oral implants: one-year results from a prospective case series.2. The-unite fixed dental prosthesis (FDP) reconstruction. *J Clin Peridontol.* 2013;40(5):553-62. <https://doi.org/10.1111/jcpe.12093> PMID:23506654.

**Cite this article :** Kokovic V, Tattan M.A Six-Month Clinical Study to Evaluate the Effects of Sodium Tripolyphosphate and Tetrapotassium Pyrophosphate Based Calculus Dissolution Oral Rinse in Patients with Zirconium Dioxide and Titanium Dental Implants *BEMS Reports*, 2016;2(2):64-70.